

Cancellation theorem

Vladimir Voevodsky

January 2002

Contents

1	Introduction	1
2	Finite correspondences	2
3	Intersecting relative cycles with divisors	5
4	Cancellation theorem	9

1 Introduction

Let $SmCor(k)$ be the category of smooth finite correspondences over a field k . Denote by $\mathbf{G}_m$ the scheme $\mathbf{A}^1 - \{0\}$. One defines the sheaf with transfers S_t^1 by the condition that $\mathbf{Z}_{tr}(\mathbf{G}_m) = S_t^1 \oplus \mathbf{Z}$ where $\mathbf{Z}$ is split off by the projection to the point and the point 1. For any scheme X consider the sheaf with transfers

$$F_X : Y \mapsto Hom(S_t^1 \otimes \mathbf{Z}_{tr}(Y), S_t^1 \otimes \mathbf{Z}_{tr}(X))$$

We prove that the obvious map $\mathbf{Z}_{tr}(X) \rightarrow F_X$ defines a quasi-isomorphism of singular simplicial complexes

$$C_*(\mathbf{Z}_{tr}(X)) \rightarrow C_*(F_X)$$

as complexes of presheaves i.e. for any Y the map of complexes of abelian groups

$$C_*(\mathbf{Z}_{tr}(X))(Y) \rightarrow C_*(F_X)(Y)$$

is a quasi-isomorphism. We then deduce from this result the cancellation theorem which asserts that if k is a perfect field then for any K, L in $DM_-^{eff}(k)$ the map

$$Hom(K, K') \rightarrow Hom(K(1), K'(1))$$

is bijective.

This result was previously known in two particular situations. For varieties over a field k with resolution of singularities it was proved in [6]. For

K' being the motivic complex $\mathbf{Z}(n)[m]$ and any field k it was proved in [4]. Both proofs are very long.

The main part of our argument does not use the assumption that we work with smooth schemes over a field and we give it for separated schemes of finite type over a noetherian base. To be able to do it we define in the first section the category of finite correspondences for separated schemes of finite type over a base. The definition is a straightforward generalization of the definition for schemes over a field based on the constructions of [3] and can be skipped. In the second section we define intersection of relative cycles with Cartier divisors and prove the properties of this construction which we need. In the third we prove our main theorem 4.6 and deduce from it the cancellation theorem over perfect fields 4.10.

In this paper we say “a relative cycle” instead of “an equidimensional relative cycle”. All schemes are separated. The letter S is typically reserved for the base scheme which is assumed to be noetherian. All the standard schemes $\mathbf{P}^1$, $\mathbf{A}^1$ etc. are over S . When no confusion is possible we write XY instead of $X \times_S Y$.

2 Finite correspondences

For a scheme X of finite type over a noetherian scheme S we denote by $c(X/S)$ the group of finite relative cycles on X over S . In [3] this group was denoted by $c_{\text{equi}}(X/S, 0)$. If S is regular or if S is normal and the characteristic of X is zero, $c(X/S)$ is the free abelian group generated by closed irreducible subsets of X which are finite over S and surjective over a connected component of S . For the general definition see [3, after Lemma 3.3.9]. A morphism $f : S' \rightarrow S$ defines the pull-back homomorphism $c(X/S) \rightarrow c(XS'/S')$ which we denote by $\text{cycl}(f)$.

For two schemes X, Y of finite type over S we define the group $c(X, Y)$ of *finite correspondences* from X to Y as $c(XY/X)$.

Let us recall the following construction from [3, §3.7]. Let $X' \rightarrow X \rightarrow S$ be morphisms of finite type, $\mathcal{W}$ a relative cycle on X' over X and $\mathcal{Z}$ a relative cycle on X over S . Then one defines a cycle $\text{Cor}(\mathcal{W}, \mathcal{Z})$ on X' as follows. Let Z_i be the components of the support of $\mathcal{Z}$ present with multiplicities n_i and $e_i : Z_i \rightarrow X$ the corresponding closed embeddings. Let $e'_i : Z_i \times_X X' \rightarrow X'$ denote the projections. We set

$$\text{Cor}(\mathcal{W}, \mathcal{Z}) = \sum_i n_i (e'_i)_* \text{cycl}(e_i)(\mathcal{W})$$

where $(e'_i)_*$ is the (proper) push-forward on cycles.

Let X, Y be schemes of finite type over S and

$$f \in c(X, Y) = c(XY/X)$$

$$g \in c(Y, Z) = c(YZ/Y)$$

finite correspondences. Let

$$p_X : XY \rightarrow Y$$

$$p_Y : XYZ \rightarrow XZ$$

be the projections. We define the composition $g \circ f$ by the formula:

$$g \circ f = (p_Y)_* \text{Cor}(\text{cycl}(p_X)(g), f) \quad (2.1)$$

This operation is linear in both arguments and thus defines a homomorphism of abelian groups

$$c(X, Y) \otimes c(Y, Z) \rightarrow c(X, Z)$$

The lemma below follows immediately from the definition of $\text{Cor}(-, -)$ and the fact that the (proper) push-forward commutes with the $\text{cycl}(-)$ homomorphisms ([3, Prop. 3.6.2]).

Lemma 2.1 *Let $Y \rightarrow X \rightarrow S$ be a sequence of morphisms of finite type, $p : Y \rightarrow Y'$ a morphism over X , $\mathcal{Y} \in \text{Cycl}(Y/X, r) \otimes \mathbf{Q}$ and $\mathcal{X} \in \text{Cycl}(X/S, s) \otimes \mathbf{Q}$. Assume that p is proper on the support of $\mathcal{Y}$. Then*

$$p_* \text{Cor}(\mathcal{Y}, \mathcal{X}) = \text{Cor}(p_*(\mathcal{Y}), \mathcal{X}).$$

Lemma 2.2 *For any $f \in c(X, Y)$, $g \in c(Y, Z)$, $h \in c(Z, T)$ one has*

$$(h \circ g) \circ f = h \circ (g \circ f).$$

Proof: Consider the following diagram

$$\begin{array}{ccccccc}
& XT & \xleftarrow{4} & XYT & \longrightarrow & YT & \\
& \uparrow 7 & & \uparrow 8 & & \uparrow 2 & \\
XZT & \xleftarrow{9} & XYZT & \longrightarrow & YZT & \longrightarrow & ZT \longrightarrow T \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
XZ & \xleftarrow{5} & XYZ & \xrightarrow{9} & YZ & \xrightarrow{1} & Z \\
& & \downarrow & & \downarrow & & \\
& & XY & \xrightarrow{3} & Y & & \\
& & \downarrow & & & & \\
& & X & & & &
\end{array}$$

where the morphisms are the obvious projections. Note that all the squares are cartesian. We will also use the projection $6 : XZ \rightarrow Z$.

We have $f \in c(XY/X)$, $g \in c(YZ/Y)$ and $h \in c(ZT/Z)$. The compositions are given by:

$$g \circ f = 5_* \text{Cor}(\text{cycl}(3)(g), f)$$

$$h \circ g = 2_* \text{Cor}(\text{cycl}(1)(h), g)$$

$$(h \circ g) \circ f = 4_* \text{Cor}(\text{cycl}(3)(h \circ g), f) = 4_* \text{Cor}(\text{cycl}(3)(2_* \text{Cor}(\text{cycl}(1)(h), g)), f)$$

$$h \circ (g \circ f) = 7_* \text{Cor}(\text{cycl}(6)(h), g \circ f) = 7_* \text{Cor}(\text{cycl}(6)(h), 5_* \text{Cor}(\text{cycl}(3)(g), f))$$

We have:

$$\begin{aligned}
& 4_* \text{Cor}(\text{cycl}(3)(2_* \text{Cor}(\text{cycl}(1)(h), g)), f) = \\
& = 4_* \text{Cor}(8_* \text{cycl}(3) \text{Cor}(\text{cycl}(1)(h), g), f) = \\
& = 4_* 8_* \text{Cor}(\text{cycl}(3) \text{Cor}(\text{cycl}(1)(h), g), f) = \\
& = 4_* 8_* \text{Cor}(\text{Cor}(\text{cycl}(1 \circ 9)(h), \text{cycl}(3)(g)), f)
\end{aligned}$$

where the first equality holds by [3, Prop. 3.6.2], the second by Lemma 2.1 and the third by [3, Th. 3.7.3]. We also have:

$$\begin{aligned}
& 7_* \text{Cor}(\text{cycl}(6)(h), 5_* \text{Cor}(\text{cycl}(3)(g), f)) = \\
& = 7_* 9_* \text{Cor}(\text{cycl}(6 \circ 5)(h), \text{Cor}(\text{cycl}(3)(g), f))
\end{aligned}$$

by [3, Lemma 3.7.1]. We conclude that $(h \circ g) \circ f = h \circ (g \circ f)$ by [3, Prop. 3.7.7].

We denote by $Cor(S)$ the category of finite correspondences whose objects are schemes of finite type over S , morphisms are finite correspondences and the composition of morphisms is defined by (2.1).

For a morphism of schemes $f : X \rightarrow Y$ let Γ_f be its graph considered as an element of $c(XY/X)$. One verifies easily that $\Gamma_{gf} = \Gamma_g \circ \Gamma_f$ and we get a functor $Sch/S \rightarrow Cor(S)$. Below we use the same symbol for a morphism of schemes and its graph considered as a finite correspondence.

The external product of cycles defines pairings

$$c(X, Y) \otimes c(X', Y') \rightarrow c(XX', YY')$$

and one verifies easily using the results of [3] that this pairing extends to a tensor structure on $Cor(S)$ with $X \otimes Y := XY$.

3 Intersecting relative cycles with divisors

Let X be a noetherian scheme and D a Cartier divisor on X i.e. a global section of the sheaf $\mathcal{M}^*/\mathcal{O}^*$. One defines the cycle $cycl(D)$ associated with D as follows. Let U_i be an open covering of X such that D_{U_i} is of the form $f_{i,+}/f_{i,-} \in \mathcal{M}^*(U_i)$. Then $cycl(D)$ is determined by the property that

$$cycl(D)|_{U_i} = cycl(f_{i,+}^{-1}(0)) - cycl(f_{i,-}^{-1}(0))$$

where on the right hand side one considers the cycles associated with closed subschemes ([3,]). One defines the support of D as the closed subset $supp(D) := supp(cycl(D))$.

We say that a cycle $\mathcal{Z} = \sum n_i z_i$ on X intersects D properly if the points z_i do not belong to $supp(D)$. Let Z_i be the closure of z_i considered as a reduced closed subscheme and $e_i : Z_i \rightarrow X$ the closed embedding. If $\mathcal{Z}$ and D intersect properly we define their intersection $(\mathcal{Z}, D)$ as the cycle

$$(\mathcal{Z}, D) := \sum n_i (e_i)_* (cycl(e_i^*(D)))$$

If $p : X \rightarrow S$ is a morphism of finite type and $\mathcal{Z}$ is a relative cycle of relative dimension d over S , we say that D intersects $\mathcal{Z}$ properly relative to p (or properly over S) if the dimension of fibers of $supp(D) \cap supp(\mathcal{Z})$ over S is $\leq d - 1$. This clearly implies that $\mathcal{Z}$ intersects D properly and $(\mathcal{Z}, D)$ is defined.

Proposition 3.1 *Assume that S is reduced and D and $\mathcal{Z}$ intersect properly over S . Then:*

1. $(\mathcal{Z}, D)$ is a relative cycle of relative dimension $d - 1$ over S
2. let $p : S' \rightarrow S$ be a morphism and $p' : XS' \rightarrow X$ the projection, assume that $(p')^*(D)$ is defined, then one has

$$p^*(\text{cycl}(Z), D) = (p^*(\text{cycl}(Z)), (p')^*(D)). \quad (3.1)$$

Proof: One can easily see that D intersects $\mathcal{Z}$ properly relative to p if and only if for any i and any point x of X there exists a neighborhood U of x such that $D|_U = f_+/f_-$ where f_+ and f_- are not zero divisors on the fibers of the maps $Z_i \rightarrow S$. Since the problem is local in X we may assume that $D = D(f)$ where f is a regular function on X which is not a zero divisor on fibers of the maps $Z_i \rightarrow S$. We write $(\mathcal{Z}, f)$ instead of $(\mathcal{Z}, D(f))$.

To prove both statements of the proposition it is clearly sufficient to show that for any fat point $\text{Spec}(k) \xrightarrow{x_0} \text{Spec}(R) \xrightarrow{x_1} S$ one has

$$(x_0, x_1)^*((\mathcal{Z}, f)) = ((x_0, x_1)^*(\mathcal{Z}), f x'_1 x'_0)$$

where x'_0, x'_1 are the projections

$$X \text{Spec}(k) \xrightarrow{x'_0} X \text{Spec}(R) \xrightarrow{x'_1} X$$

Let $j : \text{Spec}(K(R)) \rightarrow \text{Spec}(R)$ be the generic point of R and

$$pr : \text{Spec}(K(R))X \rightarrow X$$

the projection. Since S is reduced, Z_i are flat over a dense open subset of S and $(x_1 j)^*((\mathcal{Z}, f)) = \sum m_j w_j$ is defined and by Lemma 3.3 we have

$$(x_1 j)^*((\mathcal{Z}, f)) = ((x_1 j)^*(\mathcal{Z}), f \circ pr)$$

For a cycle $\mathcal{W} = \sum m_j w_j$ over the generic point of $\text{Spec}(R)$ denote by $cl \mathcal{W}$ the same cycle considered over $\text{Spec}(R)$. Since $cl(x_1 j)^*(\mathcal{Z})$ is a formal linear combination of cycles of flat subschemes, Lemma 3.3 applies and we conclude that:

$$\begin{aligned} (x_0, x_1)^*((\mathcal{Z}, f)) &= x_0^*(cl(x_1 j)^*((\mathcal{Z}, f))) = x_0^*(cl(x_1 j)^*(\mathcal{Z}), f x'_1) = \\ &= (x_0^* cl(x_1 j)^*(\mathcal{Z}), f x'_1 x'_0) = ((x_0, x_1)^*(\mathcal{Z}), f x'_1 x'_0) \end{aligned}$$

Lemma 3.2 *Let Z be a noetherian scheme and f a regular function on Z which is not a zero divisor. Let further $e_i : Z_i \rightarrow Z$ be the connected components of Z and n_i their multiplicities. Then one has:*

$$\text{cycl}(f^{-1}(0)) = \sum n_i(e_i)_*(\text{cycl}(f|_{Z_i}^{-1}(0))) \quad (3.2)$$

Proof: (P.Deligne) Replacing Z by the local scheme of a generic point of $f^{-1}(0)$ we may assume that $Z = \text{Spec}(A)$ is local of dimension 1. By definition of the cycle associated to a closed subscheme (3.2) is equivalent to

$$l(A/(f)) = \sum l(A/p_i)l(A/(p_i + (f)))$$

where p_i are the minimal ideals of A and $l(M)$ is the length of an A -module. The function $M \rightarrow l(M)$ extends to an additive function on the K_0 of the abelian category of A -modules of finite length. Consider the function $l_f : M \mapsto l(\text{cone}(M \xrightarrow{f} M))$. Equivalently we have $l_f(M) = l(M \otimes^L A/f)$ which shows that it is an additive function on $K_0(A - \text{mod})$. In $K_0(A - \text{mod})$ one has

$$A = (\sum l(A/p_i)A/p_i) + M$$

where M is a module of finite length. This implies:

$$l(A/f) = l_f(A) = \sum l(A/p_i)l_f(A/p_i) = \sum l(A/p_i)l(A/(p_i + (f))).$$

Lemma 3.3 *Let Z be a closed subscheme of X which is flat and equidimensional over S and f be a regular function on Z such that $f^{-1}(0) \cap Z$ is equidimensional over S . Then $(\text{cycl}(Z), f)$ is a relative cycle and for any morphism $p : S' \rightarrow S$ one has:*

$$p^*(\text{cycl}(Z), f) = (p^*(\text{cycl}(Z)), fp')$$

where $p' : XS' \rightarrow X$.

Proof: Lemma 3.2 implies that

$$(\text{cycl}(Z), f) = \text{cycl}(f|_Z^{-1}(0)).$$

Our condition on f implies that, for any $s \in S$, f restricted to the fiber Z_s is not a zero divisor. Since Z is flat over S this implies that $f|_Z^{-1}(0)$ is flat

over S (see e.g. [1,]). Since Z is equidimensional of relative dimension d , $f|_Z^{-1}(0)$ is equidimensional of relative dimension $d - 1$. We conclude by [3,] that $\text{cycl}(f|_Z^{-1}(0))$ is a relative cycle.

For $p; S' \rightarrow S$ we have:

$$\begin{aligned} p^*(\text{cycl}(Z), f) &= p^*(\text{cycl}(f|_Z^{-1}(0))) = \text{cycl}((fp')|_{ZS'}^{-1}(0)) = \\ &= (\text{cycl}(ZS'), fp') = (p^*(\text{cycl}(Z)), fp'). \end{aligned}$$

Corollary 3.4 *Let $X' \xrightarrow{f} X \rightarrow S$ be morphisms of finite type, $\mathcal{Z}$ a relative cycle on X over S and $\mathcal{W}$ a relative cycle on X' over X of dimension 0. Let further D be a Cartier divisor on X' which intersects $\mathcal{W}$ properly over X . Then D intersects $\text{Cor}(\mathcal{W}, \mathcal{Z})$ properly over S and one has:*

$$(\text{Cor}(\mathcal{W}, \mathcal{Z}), D) = \text{Cor}((\mathcal{W}, D), \mathcal{Z}) \quad (3.3)$$

Proof: It is a straightforward corollary of the definition of $\text{Cor}(-, -)$ and (3.1).

Lemma 3.5 *Let $f : X' \rightarrow X$ be a morphism of schemes of finite type over S , $\mathcal{Z}$ a relative cycle on X' such that f is proper on $\text{supp}(\mathcal{Z})$ and D a Cartier divisor on X . Assume that $f^*(D)$ is defined and $\mathcal{Z}$ intersects $f^*(D)$ properly over S . Then $f_*(\mathcal{Z})$ intersects D properly over S and one has:*

$$f_*(\mathcal{Z}, f^*(D)) = (f_*(\mathcal{Z}), D) \quad (3.4)$$

Proof: Let d be the relative dimension of $\mathcal{Z}$ over S . To see that $f_*(\mathcal{Z})$ intersects D properly over S we need to check that the dimension of the fibers of $\text{supp}(D) \cap \text{supp}(f_*(\mathcal{Z}))$ over S is $\leq d - 1$. This follows from our assumption and the inclusion

$$\begin{aligned} \text{supp}(D) \cap \text{supp}(f_*(\mathcal{Z})) &\subset \text{supp}(D) \cap f(\text{supp}(\mathcal{Z})) = \\ &= f(f^{-1}(\text{supp}(D)) \cap \text{supp}(\mathcal{Z})) = f(\text{supp}(f^*(D)) \cap \text{supp}(\mathcal{Z})) \end{aligned}$$

To verify (3.4) it is sufficient to consider the situation locally around the generic points of $f(\text{supp}(f^*(D)) \cap \text{supp}(\mathcal{Z}))$. Therefore we may assume that

$D = D(g)$ is the divisor of a regular function g and $\mathcal{Z} = z$ is just one point with the closure Z . Replacing X' by Z and X by $f(Z)$ we may assume that X, X' are integral, f is surjective and X is local of dimension 1. Let $A = \mathcal{O}(X)$, $B = \mathcal{O}(X')$. As in the proof of 3.2, consider the function $l_g : M \mapsto l_A(M \otimes^L A/g)$ on $K_0(A\text{-mod})$. This function vanishes on modules with the support in the closed point which implies that

$$l_g(B) = \deg(f)l_g(A) = \deg(f)l_A(A/g)$$

On the other hand $l_g(A) = l_A(B/(f^*(g)))$. Let x'_i be the closed points of X' , k'_i their residue fields and k the residue field of the closed point of X . Let further M_i be the part of $B/(f^*(g))$ supported in x'_i . One can easily see that $l_A(B/(f^*(g))) = \sum_i [k'_i : k]l_B(M_i)$. Combining our equalities we get:

$$\deg(f)l_A(A/g) = \sum_i [k'_i : k]l_B(M_i) \quad (3.5)$$

which is equivalent to (3.4).

4 Cancellation theorem

Consider a finite correspondence

$$\mathcal{Z} \in c(\mathbf{G}_m X, \mathbf{G}_m Y) = c(\mathbf{G}_m X \mathbf{G}_m Y / \mathbf{G}_m X).$$

Let f_1, f_2 be the projections to the first and the second copy of $\mathbf{G}_m$ respectively and let g_n denote the rational function $(f_1^{n+1} - 1)/(f_1^{n+1} - f_2)$ on $\mathbf{G}_m X \mathbf{G}_m Y$.

Lemma 4.1 *For any $\mathcal{Z}$ there exists N such that for all $n \geq N$ the divisor of g_n intersects $\mathcal{Z}$ properly over X and the cycle $(\mathcal{Z}, D(g_n))$ is finite over X .*

Proof: Let $\bar{f}_1 \times \bar{q} : \bar{C} \rightarrow \mathbf{P}^1 X$ be a finite morphism which extends the projection $\text{supp}(\mathcal{Z}) \rightarrow \mathbf{G}_m X$. Let N be an integer such that the rational function $\bar{f}_1^N / f_2$ is regular in a neighborhood of $\bar{f}_1^{-1}(0)$ and the rational function $f_2 / \bar{f}_1^N$ is regular in a neighborhood of $\bar{f}_1^{-1}(\infty)$. Then for any $n \geq N$ one has:

1. the restriction of $g_n f_2$ to $\text{supp}(\mathcal{Z})$ is regular on a neighborhood of $\bar{f}_1^{-1}(0)$ and equals 1 on $\bar{f}_1^{-1}(0)$
2. the restriction of g_n to $\text{supp}(\mathcal{Z})$ is regular a neighborhood of $\bar{f}_1^{-1}(\infty)$ and equals 1 on $\bar{f}_1^{-1}(\infty)$

Conditions (1),(2) imply that the divisor of g_n intersects $\mathcal{Z}$ properly over X and that the relative cycle $(\mathcal{Z}, D(g_n))$ is finite over X .

If $(\mathcal{Z}, D(g_n))$ is defined as a finite relative cycle we let $\rho_n(\mathcal{Z}) \in c(X, Y)$ denote the projection of $(\mathcal{Z}, D(g_n))$ to XY .

Remark 4.2 Note that we can define a finite correspondence $\rho_g(\mathcal{Z}) : X \rightarrow Y$ for any function g satisfying the conditions (1),(2) in the same way as we defined $\rho_n = \rho_{g_n}$. In particular, if n and m are large enough then the function $tg_n + (1-t)g_m$ defines a finite correspondence $h = h_{n,m} : X\mathbf{A}^1 \rightarrow Y$ such that $h|_{X \times \{0\}} = \rho_m(\mathcal{Z})$ and $h|_{X \times \{1\}} = \rho_n(\mathcal{Z})$, i.e. we get a canonical $\mathbf{A}^1$ -homotopy from $\rho_m(\mathcal{Z})$ to $\rho_n(\mathcal{Z})$.

Lemma 4.3 *[(i)]*

For a finite correspondence $\mathcal{W} : X \rightarrow Y$ and any $n \geq 1$ one has $\rho_n(\text{Id}_{\mathbf{G}_m} \otimes \mathcal{W}) = \mathcal{W}$

2. Let e_X be the composition $\mathbf{G}_m X \xrightarrow{pr} X \xrightarrow{\{1\} \times \text{Id}} \mathbf{G}_m X$. Then $\rho_n(e_X) = 0$ for any $n \geq 0$.

Proof: The cycle on $\mathbf{G}_m X \mathbf{G}_m Y$ over $\mathbf{G}_m X$ which represents $\text{Id}_{\mathbf{G}_m} \otimes \mathcal{W}$ is $\Delta_*(\mathbf{G}_m \times \mathcal{W})$ where Δ is the diagonal embedding $\mathbf{G}_m XY \rightarrow \mathbf{G}_m X \mathbf{G}_m Y$. The cycle $(\Delta_*(\mathbf{G}_m \times \mathcal{W}), g_n)$ is $\Delta_*(D \otimes \mathcal{W})$ where D is the divisor of the function $(t^{n+1} - 1)/(t^{n+1} - t)$ on $\mathbf{G}_m$. The push-forward of $\Delta_*(D \otimes \mathcal{W})$ to XY is the cycle $\deg(D)\mathcal{W}$. Since $\deg(D) = 1$ we get the first statement of the lemma.

The cycle $\mathcal{Z}$ on $\mathbf{G}_m X \mathbf{G}_m X$ representing e_X is the image of the embedding $\mathbf{G}_m X \rightarrow \mathbf{G}_m X \mathbf{G}_m X$ which is diagonal on X and of the form $t \mapsto (t, 1)$ on $\mathbf{G}_m$. This shows that the restriction of g_n to $\text{supp}(\mathcal{Z})$ equals 1 and $(\mathcal{Z}, D(g_n)) = 0$.

Lemma 4.4 *Let $\mathcal{Z} : \mathbf{G}_m X \rightarrow \mathbf{G}_m Y$ be a finite correspondence such that $\rho_n(\mathcal{Z})$ is defined. Then for any finite correspondence $\mathcal{W} : X' \rightarrow X$, $\rho_n(\mathcal{Z} \circ (\text{Id}_{\mathbf{G}_m} \otimes \mathcal{W}))$ is defined and one has*

$$\rho_n(\mathcal{Z} \circ (\text{Id}_{\mathbf{G}_m} \otimes \mathcal{W})) = \rho_n(\mathcal{Z}) \circ \mathcal{W} \quad (4.1)$$

Proof: Let us show that (4.1) holds. In the process it will become clear that the left hand side is defined. We can write $\rho_n(\mathcal{Z}) \circ \mathcal{W}$ as the composition

$$X' \xrightarrow{\mathcal{W}} X \xrightarrow{(\mathcal{Z}, D(g_n))} \mathbf{G}_m \mathbf{G}_m Y \xrightarrow{pr} Y$$

and $\rho_n(\mathcal{Z} \circ (Id_{\mathbf{G}_m} \otimes \mathcal{W}))$ as the composition

$$X' \xrightarrow{\mathcal{Y}} \mathbf{G}_m \mathbf{G}_m Y \xrightarrow{pr} Y$$

where $\mathcal{Y} = (\mathcal{Z} \circ (Id_{\mathbf{G}_m} \otimes \mathcal{W}), D(g_n))$. Consider the diagram

$$\begin{array}{ccccc} \mathbf{G}_m X' \mathbf{G}_m Y & \xleftarrow{p_1} & \mathbf{G}_m X' X \mathbf{G}_m Y & \longrightarrow & \mathbf{G}_m X \mathbf{G}_m Y \\ & & \downarrow & & \downarrow \\ & & X' X & \xrightarrow{p_2} & X \\ & & \downarrow & & \\ & & X' & & \end{array}$$

where the arrows are the obvious projections. If we consider $\mathcal{Z}$ as a cycle of dimension 1 over X then the cycle $\mathcal{Z} \circ (Id_{\mathbf{G}_m} \otimes \mathcal{W})$, considered as a cycle over X' , is $(p_1)_* Cor(cycl(p_2)(\mathcal{Z}), \mathcal{W})$ and we have

$$\begin{aligned} & ((p_1)_* Cor(cycl(p_2)(\mathcal{Z}), \mathcal{W}), D(g_n)) = \\ & = (p_1)_*(Cor(cycl(p_2)(\mathcal{Z}), \mathcal{W}), D(g_n)) = (p_1)_* Cor((cycl(p_2)(\mathcal{Z}), D(g_n)), \mathcal{W}) = \\ & = (p_1)_* Cor(cycl(p_2)(\mathcal{Z}, D(g_n)), \mathcal{W}) \end{aligned}$$

where the first equality holds by (3.4), the second by (3.3) and the third by (3.1).

The last expression represents the composition $\mathcal{W} \circ (\mathcal{Z}, D(g_n))$ and we conclude that

$$\rho_n(\mathcal{Z}) \circ \mathcal{W} = \rho_n(\mathcal{Z} \circ (Id_{\mathbf{G}_m} \otimes \mathcal{W}))$$

Lemma 4.5 *Let $\mathcal{Z} : \mathbf{G}_m X \rightarrow \mathbf{G}_m Y$ be a finite correspondence such that $\rho_n(\mathcal{Z})$ is defined. Then for any morphism of schemes $f : X' \rightarrow Y'$, $\rho_n(\mathcal{Z} \otimes f)$ is defined and one has*

$$\rho_n(\mathcal{Z} \otimes f) = \rho_n(\mathcal{Z}) \otimes f \tag{4.2}$$

Proof: Consider the diagram

$$\begin{array}{ccccc}
\mathbf{G}_m X X' \mathbf{G}_m Y Y' & \xleftarrow{p_1} & \mathbf{G}_m X X' \mathbf{G}_m Y & \longrightarrow & \mathbf{G}_m X \mathbf{G}_m Y \\
& & \downarrow & & \downarrow \\
& & X X' & \xrightarrow{p_2} & X
\end{array}$$

where p_1 is defined by the embedding $X' \xrightarrow{f \times Id} X'Y'$ and the rest of the morphisms are the obvious projections. Consider $\mathcal{Z}$ as a cycle over X . Then $\rho_n(\mathcal{Z} \otimes f)$ is given by the composition

$$\mathbf{G}_m X X' \xrightarrow{\mathcal{Y}_1} \mathbf{G}_m \mathbf{G}_m Y \xrightarrow{pr} Y Y'$$

where $\mathcal{Y}_1 = ((p_1)_* cycl(p_2)(\mathcal{Z}), g_n)$ and $\rho_n(\mathcal{Z}) \otimes f$ by the composition

$$\mathbf{G}_m X X' \xrightarrow{\mathcal{Y}_2} \mathbf{G}_m \mathbf{G}_m Y \xrightarrow{pr} Y Y'$$

where $\mathcal{Y}_2 = (p_1)_*(cycl(p_2)((\mathcal{Z}, g_n)))$. The equality $\mathcal{Y}_1 = \mathcal{Y}_2$ follows from (3.4) and (3.1).

For our next result we need to use presheaves with transfers. A presheaf with transfers on Sch/S is an additive contravariant functor from $Cor(S)$ to the category of abelian groups. For X in Sch/S we let $\mathbf{Z}_{tr}(X)$ denote the functor represented by X on $Cor(S)$. One defines tensor product of presheaves with transfers in the usual way such that $\mathbf{Z}_{tr}(X) \otimes \mathbf{Z}_{tr}(Y) = \mathbf{Z}_{tr}(X \times Y)$. To simplify notations we will write X instead of $\mathbf{Z}_{tr}(X)$ and identify morphisms $\mathbf{Z}_{tr}(X) \rightarrow \mathbf{Z}_{tr}(Y)$ with finite correspondences $X \rightarrow Y$. Note in particular that $\mathbf{G}_m$ denotes the presheaf with transfers $\mathbf{Z}_{tr}(\mathbf{G}_m)$ not the presheaf with transfers represented by $\mathbf{G}_m$ as a scheme. To preserve compatibility with the notation XY for the product of X and Y we write FG for the tensor product of presheaves with transfers F and G .

Let S_t^1 denote the presheaf with transfers $ker(\mathbf{G}_m \rightarrow S)$. We consider it as a direct summand of $\mathbf{G}_m$ with respect to the projection $Id - e$ where e is defined by the composition $\mathbf{G}_m \rightarrow S \xrightarrow{1} \mathbf{G}_m$. In the following theorem we let $f \cong g$ denote that the morphisms f and g are $\mathbf{A}^1$ -homotopic.

Theorem 4.6 *Let F be a presheaf with transfers such that there is an epimorphism $X \rightarrow F$ for a scheme X . Let $\phi : S_t^1 \otimes F \rightarrow S_t^1 Y$ be a morphism. Then there exists a unique up to an $\mathbf{A}^1$ -homotopy morphism $\rho(\phi) : F \rightarrow Y$ such that $Id_{S_t^1} \otimes \rho(\phi) \cong \phi$.*

Proof: Let us fix an epimorphism $p : X \rightarrow F$. Then the morphism ϕ defines a finite correspondence $\mathcal{Z} : \mathbf{G}_m X \rightarrow \mathbf{G}_m Y$ and for n sufficiently large we may consider $\rho_n(\mathcal{Z}) : X \rightarrow Y$. Lemma 4.4 implies immediately that $\rho_n(\mathcal{Z})$ vanishes on $\ker(p)$ and therefore it defines a morphism $\rho_n(\phi) : F \rightarrow X$.

Consider a morphism ϕ of the form $Id_{S_t^1} \otimes \psi$. Then $\mathcal{Z}$ is of the form $(Id_{\mathbf{G}_m} - e) \otimes \mathcal{W}$ where $\mathcal{W} : X \rightarrow Y$ corresponds to ψ . By Lemma 4.3 we have $\rho_n(\mathcal{Z}) = \mathcal{W}$ and therefore $\rho_n(Id_{S_t^1} \otimes \psi) = \psi$ for any $n \geq 1$. If ρ, ρ' are two morphisms such that $Id_{S_t^1} \otimes \rho \cong \phi$ and $Id_{S_t^1} \otimes \rho' \cong \phi$ then for a sufficiently large n we have

$$\rho = \rho_n(Id_{S_t^1} \otimes \rho) \cong \rho_n(Id_{S_t^1} \otimes \rho') = \rho'$$

This implies the uniqueness part of the theorem.

To prove the existence let us show that for a sufficiently large n one has $Id_{S_t^1} \otimes \rho_n(\phi) \cong \phi$. Let $\tilde{\phi}$ be the morphism $\mathbf{G}_m F \rightarrow \mathbf{G}_m Y$ defined by ϕ and let

$$\tilde{\phi}^* : F\mathbf{G}_m \rightarrow Y\mathbf{G}_m$$

be the morphism obtained from $\tilde{\phi}$ by the obvious permutation.

Lemma 4.7 *The morphisms $\tilde{\phi} \otimes (Id_{\mathbf{G}_m} - e)$ and $(Id_{\mathbf{G}_m} - e) \otimes \tilde{\phi}^*$ are $\mathbf{A}^1$ -homotopic.*

Proof: One can easily see that these two morphisms are obtained from the morphisms

$$\phi \otimes Id_{S_t^1}, Id_{S_t^1} \otimes \phi^* : S_t^1 F S_t^1 \rightarrow S_t^1 Y S_t^1$$

by using the standard direct sum decomposition. One can see further that $\phi \otimes Id_{S_t^1} = \sigma_Y(Id_{S_t^1} \otimes \phi^*)\sigma_F$ where σ_F and σ_Y are the permutations of the two copies of S_t^1 in $S_t^1 F S_t^1$ and $S_t^1 Y S_t^1$ respectively. Lemma 4.8 below implies now that $\phi \otimes Id_{S_t^1} \cong Id_{S_t^1} \otimes \phi^*$.

Lemma 4.8 *The permutation on $S_t^1 S_t^1$ is $\mathbf{A}^1$ -homotopic to $\{-1\}Id \otimes Id$ where $\{-1\} : S_t^1 \rightarrow S_t^1$ is defined by the morphism $\mathbf{G}_m \xrightarrow{x \mapsto x^{-1}} \mathbf{G}_m$.*

Proof: The same arguments as the ones used in [2, p.142] show that for any scheme X and any pair of invertible functions f, g on X the morphism $X \xrightarrow{f \otimes g} S_t^1 S_t^1$ is $\mathbf{A}^1$ -homotopic to the morphism $g \otimes f^{-1}$. This implies immediately that the permutation on $S_t^1 S_t^1$ is $\mathbf{A}^1$ -homotopic to the morphism

$Id \otimes (\{-1\}Id)$ where $\{-1\}Id : S_t^1 \rightarrow S_t^1$ is the morphism defined by the map $\mathbf{G}_m \xrightarrow{x \mapsto x^{-1}} \mathbf{G}_m$.

For a sufficiently large n we have

$$\rho_n(\phi \otimes (Id_{\mathbf{G}_m} - e)) = \rho_n(\phi) \otimes (Id_{\mathbf{G}_m} - e)$$

by Lemma 4.5. On the other hand

$$\rho_n((Id_{\mathbf{G}_m} - e) \otimes \phi^*) = \phi^*$$

by Lemma 4.3. By Lemma 4.7 we conclude that

$$\phi^* \cong \rho_n(\phi) \otimes (Id_{\mathbf{G}_m} - e)$$

which is equivalent to $Id_{S_t^1} \otimes \rho_n(\phi) \cong \phi$. Theorem 4.6 is proved.

Corollary 4.9 *Denote by F_Y the presheaf*

$$X \mapsto Hom(S_t^1 X, S_t^1 Y)$$

and consider the obvious map $Y \rightarrow F_Y$. Then for any X the corresponding map of complexes of abelian groups

$$C_*(Y)(X) \rightarrow C_*(F_Y)(X)$$

is a quasi-isomorphism

Proof: Let $\Delta^n \cong \mathbf{A}^n$ be the standard algebraic simplex and $\partial\Delta^n$ the subpresheaf in Δ^n which is the union of the images of the face maps $\Delta^{n-1} \rightarrow \Delta^n$. Then the n -th homology group of the complex $C_*(F)(X)$ for any F is the group of homotopy classes of maps from $X \otimes (\Delta^n / \partial\Delta^n)$ to F . Our result now follows directly from 4.6.

Corollary 4.10 *Let k be a perfect field. Then for any K, L in $DM_-^{eff}(k)$ the map $Hom(K, L) \rightarrow Hom(K(1), L(1))$ is a bijection.*

Proof: Since DM_-^{eff} is generated by objects of the form X it is enough to check that for smooth schemes X, Y over k and $n \in \mathbf{Z}$ one has

$$Hom(S_t^1 X, S_t^1 Y[n]) = Hom(X, Y[n])$$

By Corollary 4.9 we know that the map

$$Y \rightarrow F_Y = \underline{Hom}(S_t^1, S_t^1 Y)$$

is an isomorphism in DM . Let us show now that for any sheaf with transfers F and any X one has

$$Hom_{DM}(S_t^1 X, F[n]) = Hom_{DM}(X, \underline{Hom}(S_t^1, F)[n]) \quad (4.3)$$

The left hand side of (4.3) is the hypercohomology group $\mathbf{H}^n(\mathbf{G}_m X, C_*(F))$ modulo the subgroup $\mathbf{H}^n(X, C_*(F))$. The right hand side is the hypercohomology group $\mathbf{H}^n(X, C_* \underline{Hom}(\mathbf{G}_m, F))$ modulo similar subgroup. Let $p : \mathbf{G}_m X \rightarrow X$ be the projection. It is easy to see that (4.3) asserts that $\mathbf{R}p_*(C_*(F)) \cong C_*(p_*(F))$. There is a spectral sequence which converges to the cohomology sheaves of $\mathbf{R}p_*(C_*(F))$ and starts with the higher direct images $R^i p_*(\underline{H}^j(C_*(F)))$. We need to verify that $R^i p_*(\underline{H}^j(C_*(F))) = 0$ for $i > 0$ and that $p_*(\underline{H}^j(C_*(F))) = \underline{H}^j(C_*(p_*(F)))$. Both statements follow from [5, Prop. 4.34, p.124] and the comparison of Zariski and Nisnevich cohomology for homotopy invariant presheaves with transfers.

References

- [1] J.S. Milne. *Etale Cohomology*. Princeton Univ. Press, Princeton, NJ, 1980.
- [2] Andrei Suslin and Vladimir Voevodsky. Bloch-Kato conjecture and motivic cohomology with finite coefficients. In *The arithmetic and geometry of algebraic cycles*, pages 117–189. Kluwer, 2000.
- [3] Andrei Suslin and Vladimir Voevodsky. Relative cycles and Chow sheaves. In *Cycles, transfers and motivic homology theories*, Annals of Math Studies, pages 10–86. Princeton Univ. Press, 2000.
- [4] Vladimir Voevodsky. Motivic cohomology are isomorphic to higher Chow groups. www.math.uiuc.edu/K-theory/378, 1999.

- [5] Vladimir Voevodsky. Cohomological theory of presheaves with transfers. In *Cycles, transfers and motivic homology theories*, Annals of Math Studies, pages 87–137. Princeton Univ. Press, 2000.
- [6] Vladimir Voevodsky. Triangulated categories of motives over a field. In *Cycles, transfers and motivic homology theories*, Annals of Math Studies, pages 188–238. Princeton Univ. Press, 2000.